



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

average of the past five years being 221, past ten years 307, past fifteen years 348. There were more deaths than in August of the past two years, the distribution being general, with no special epidemic reported. No smallpox has existed in the State since July. Reports indicate an unusual prevalence of typhoid fever in all parts of the State, and give promise of an excessive mortality in the succeeding month.

*Yonkers*.—Month of August, 1900. Estimated population, 45,000. Total number of deaths, 72, including 5 from phthisis pulmonalis.

NORTH CAROLINA—*Charlotte*.—Month of September, 1900. Estimated population, 29,000. Total number of deaths, 28, including 1 from enteric fever.

*Wilmington*.—Month of September, 1900. Estimated population, 27,000. Total number of deaths, 77, including 6 from phthisis pulmonalis.

PENNSYLVANIA—*Columbia*.—Month of September, 1900. Estimated population, 12,500. Total number of deaths, 13, including 1 from diphtheria.

TENNESSEE—*Chattanooga*.—Month of September, 1900. Estimated population, 40,000—white, 27,000; colored, 13,000. Total number of deaths, 29—white, 18; colored, 11—including diphtheria, 1; enteric fever, 2, and 6 from tuberculosis.

VIRGINIA—*Lynchburg*.—Month of September, 1900. Estimated population, 30,000. Total number of deaths, 28, including diphtheria, 1; enteric fever, 4, and 5 from phthisis pulmonalis.

WISCONSIN—*Milwaukee*.—Month of August, 1900. Population, United States census of 1900, 285,315. Total number of deaths, 441, including diphtheria, 8; enteric fever, 6; measles, 2; scarlet fever, 1; whooping cough, 5, and 40 from tuberculosis.

*Report of immigrants inspected at the port of New Orleans, La., during the month of September, 1900.*

Total number of immigrants inspected, 107; number passed, 106; number certified for deportation on account of dangerous contagious or loathsome diseases, or for other physical causes, 1.

*Disposition of immigrants certified for deportation*.—Number cases pending at beginning of month, none; number cases certified for deportation during month, 1; total to be accounted for, 1; number cases deported, none; number cases admitted, none; number cases pending at close of month, 1.

C. P. WERTENBAKER,  
*Passed Assistant Surgeon, U. S. M. H. S.*